



## Hepatitis Delta virus genotype 8 infection in Northeast Brazil: Inheritance from African slaves?

L.M.F. Barros<sup>a</sup>, M.S. Gomes-Gouvêa<sup>b,\*</sup>, J.R.R. Pinho<sup>b</sup>, M.V. Alvarado-Mora<sup>b</sup>, A. Dos Santos<sup>a</sup>, M.C.J. Mendes-Corrêa<sup>c</sup>, A.J.M. Caldas<sup>a</sup>, M.T. Sousa<sup>a</sup>, M.D.C. Santos<sup>a</sup>, A.S.P. Ferreira<sup>a</sup>

<sup>a</sup> Hospital Universitário da Universidade Federal do Maranhão, São Luís, Maranhão, Brazil

<sup>b</sup> São Paulo Institute of Tropical Medicine, University of São Paulo, Brazil

<sup>c</sup> Department of Infectious Diseases, Medical School, University of São Paulo, São Paulo, Brazil

### ARTICLE INFO

#### Article history:

Received 16 May 2011

Received in revised form 10 July 2011

Accepted 11 July 2011

Available online 21 July 2011

#### Keywords:

Hepatitis Delta virus

Hepatitis B virus

HBV

HDV

Genotype

Bayesian analysis

Brazil

Prevalence

### ABSTRACT

Hepatitis Delta virus (HDV) is endemic worldwide, but its prevalence varies in different geographical areas. While in the Brazilian Amazon, HDV is known to be endemic and to represent a significant public health problem, few studies have assessed its prevalence in other regions in the country. This study evaluated the seroprevalence of HDV among HBsAg chronic carriers from Maranhão state, a region located in the Northeast of Brazil. Among 133 patients, 5 had anti-HD, of whom 3 had HDV RNA. HDV genotypes were characterized by Bayesian phylogenetic analysis of nucleotide sequences from the HDAg coding region. HDV-3 was identified in one patient who lives in Maranhão, but was born in Amazonas state (Western Amazon basin). Phylogenetic analysis shows that this HDV-3 sequence grouped with other HDV-3 sequences isolated in this state, which suggests that the patient probably contracted HDV infection there. Surprisingly, the other two patients were infected with HDV-8, an African genotype. These patients were born and have always lived in Urbano Santos, a rural county of Maranhão state, moreover they had never been to Africa and denied any contact with people from that continent. This is the first description of the HDV-8 in non-native African populations. This genotype may have been introduced to Brazil through the slaves brought to the country from the West Africa regions during the 16–18th centuries. Our results indicate that the need of clinical and epidemiological studies to investigate the presence of this infection in other areas in Brazil.

© 2011 Elsevier B.V. All rights reserved.

### 1. Introduction

Hepatitis Delta virus (HDV) is a highly pathogenic virus that causes acute and chronic liver diseases. HDV is found only in individuals positive for the hepatitis B virus (HBV) surface antigen (HBsAg) because the virus is defective and needs a helper function provided by HBsAg that is used as its own envelope protein for particle assembly and viral transmission across the human host (Wedemeyer and Manns, 2010).

The prevalence of HDV infection varies according to the geographical area. Around 15 million HDV carriers are estimated worldwide (Rizzetto, 2009). In Brazil, HDV is endemic in some areas of the Amazon region where it is associated with severe forms

of the disease (Bensabath and Dias, 1983; Bensabath et al., 1987; Gomes-Gouvea et al., 2008, 2009). In the Amazon region, anti-HD antibodies can be found in up to 34% of HBsAg carriers (Fonseca et al., 1988). Except for the studies done in this endemic region, studies on HDV prevalence in Brazil are scarce (Ferraz et al., 1985; Oliveira et al., 1999; Strauss et al., 1987).

HDV genomes isolated from around the world show 39% divergence over the entire RNA genome. This HDV diversity is currently classified into eight genotypes named HDV-1 to HDV-8. Distinct clinical courses are associated with different HDV genotypes: HDV-1 has been associated with a broad spectrum of pathogenicity, HDV-2 and HDV-4 cause milder forms of liver disease, and HDV-3 has been associated with outbreaks of fulminant hepatitis. The pathogenicities of genotypes 5–8 are not well known.

HDV-1 is geographically widespread (Shakil et al., 1997), but all other genotypes are closely associated with specific geographic areas. HDV-2 and HDV-4 are found in East Asia (Ivaniushina et al., 2001; Lee et al., 1996; Sakugawa et al., 1999); HDV-3 has been isolated in the northern area of South America only (Amazon Basin of Brazil, Peru, Colombia, and Venezuela) (Casey, 1996; Casey et al.,

\* Corresponding author at: Laboratório de Gastroenterologia e Hepatologia Tropical, Instituto de Medicina Tropical, FMUSP, Av. Dr. Enéas Carvalho de Aguiar, 500, 05403-000 São Paulo, SP, Brazil.

E-mail addresses: [gomesmic@yahoo.com.br](mailto:gomesmic@yahoo.com.br), [gomesmic@usp.br](mailto:gomesmic@usp.br) (M.S. Gomes-Gouvêa).

1993; Gomes-Gouvea et al., 2008; Quintero et al., 2001); and HDV-5 to HDV-8 have been identified in individuals from Africa (Le Gal et al., 2006; Makuwa et al., 2008, 2009; Radjef et al., 2004).

In this study we describe the presence of HDV infection in chronic HBsAg carriers from a region in Brazil other than the HDV endemic area and report for the first time the infection with HDV genotype 8 in non-native African populations.

## 2. Materials and methods

### 2.1. Patients

A total of 133 patients with chronic hepatitis B virus infection (HBsAg positive for at least 6 months) and followed at the Center for Liver Studies, University Hospital of the Federal University of Maranhão, Brazil, between January 2008 and February 2010 were enrolled in this study. Blood samples were collected and the sera separated and stored at  $-70^{\circ}\text{C}$ .

Patients' demographic and clinical data were retrieved from their medical files. All patients were from Maranhão, a state in the Northeast of Brazil. Most of them (69.2%) were from São Luís, the state capital, and the remainder were from different regions in the state (Fig. 1).

The status of chronic hepatitis B was defined according to the following criteria:

- (1) *Inactive carrier*: positive HBsAg, negative HBeAg, positive anti-HBe, normal alanine aminotransferase (ALT) levels, and HBV DNA below 2000 UI/mL.
- (2) *Chronic hepatitis B*: positive HBsAg, positive HBeAg, high ALT levels or positive HBsAg, negative HBeAg, positive anti-HBe, and HBV DNA higher than 2000 UI/mL.
- (3) *Hepatic cirrhosis*: positive HBsAg with positive or negative HBeAg regardless of the viral load, signs or symptoms of advanced hepatic disease (ascite and/or presence of esophageal varices), or histological diagnosis of hepatic cirrhosis.
- (4) *Immunotolerance*: positive HBsAg and HBeAg with normal ALT levels.

This study was approved by the local institution ethics committee and all patients provided informed consent.

### 2.2. Serological tests

All serum samples were tested for hepatitis B serological markers (HBsAg, total anti-HBc, anti-HBs, HBeAg, anti-HBe) and HDV antibody (anti-HD) using enzyme-linked immunosorbent assay (ELISA) kits (DiaSorin, Italy).

### 2.3. HDV RNA and HBV DNA amplification and sequencing

Viral nucleic acids were detected by PCR in all samples that were both positive for HBsAg and anti-HD. For this, HBV DNA and HDV RNA were initially extracted using QIAamp DNA mini kit and QIAamp Viral RNA viral mini kit (Qiagen, Germany), respectively, according to the manufacturer's instructions.

A fragment of the delta antigen genomic region (403 nucleotides) was amplified by nested polymerase chain reaction (PCR), as previously described (Gomes-Gouvea et al., 2008). HBV DNA detection was performed by a quantitative in-house real-time PCR assay (sensitivity = 50 UI/mL) (Sitnik et al., 2010). HBV genotypes were characterized by amplifying a nested PCR 1306 nucleotide fragment partially comprising HBsAg and the DNA polymerase coding regions (S/POL) using primers PS3132F/2920R and PS3201F/P1285R. To avoid false-positive results, the precau-

tions and procedures suggested by Kwok and Higuchi (1989) were strictly followed.

Amplified PCR products were purified using ChargeSwitch® PCR Clean-Up Kit (Life Technologies, USA). The sequencing reactions were carried out from both strands using inner primers and fluorescence-labeled dideoxynucleotide chain terminators using a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, USA). The sequences were determined in an automated DNA sequencer (ABI Prism 3100 Genetic Analyser, Applied Biosystems, USA). The quality of each electropherogram was evaluated using the Phred-Phrap software and consensus sequences were obtained by alignment of both sequenced strands (sense and antisense) using CAP3 software available at the web page *Electropherogram quality analysis Phred* (<http://asparagin.cenargen.embrapa.br/phph/>).

Sequences were submitted to EMBL/GenBank/DBJ under accession numbers JF298898–JF298900 (HDV sequences), and JF298901–JF298903 (HBV sequences).

### 2.4. Phylogenetic analyses

Genotyping classification for HBV and HDV was performed on the basis of phylogenetic reconstructions using representative reference sequences from different HDV genotypes ( $n = 105$ ) and HBV subgenotypes ( $n = 281$ ) obtained from the GenBank public database (<http://www.ncbi.nlm.nih.gov/>) (data available upon request). Sequences were aligned using CLUSTALX and edited using BioEdit software. Bayesian phylogenetic analyses were conducted using the Markov Chain Monte Carlo (MCMC) simulation implemented in BEAST v.1.6.1 (Drummond and Rambaut, 2007). HDV and HBV datasets were analyzed under relaxed uncorrelated lognormal molecular clock using the model of nucleotide substitution (GTR + G + I) obtained previously by Modeltest v3.7 (Posada and Crandall, 1998) and 10 million generations were sufficient to obtain the convergence of parameters. The maximum clade credibility (MCC) tree was obtained from summarizing the 10,000 substitution trees and then 10% of burn-in was removed using Tree Annotator v.1.6.1 (Drummond and Rambaut, 2007).

## 3. Results

Altogether, 133 samples from chronic HBsAg carriers were screened for the presence of antibodies against HDAg by serology. Among these patients, 69 (51.9%) were women and 64 (48.1%) were men, their mean age was 40.4 years (ranging from 13 to 84 years) and in terms of race (based on skin color), 66.2% were brown (*pardo*), 18.8% were black, and 15% were white. Sixty-nine (52%) patients were inactive carriers, 39 (29.5%) had chronic hepatitis, 15 (11%) had cirrhosis, and 10 (7.5%) were in the immunotolerance phase.

Out of 133 samples, 5 (3.8%) were positive for anti-HD and HDV RNA was detected in 3 (60%) of these anti-HD positive samples. All anti-HD positive patients showed detectable HBV viral load. HBV-DNA levels, HBeAg and anti-HBe status, HBV and HDV genotypes, and clinical/epidemiological data of anti-HD positive patients are summarized in Table 1.

A phylogenetic tree constructed with sequences of the partial delta antigen genomic region of HDV RNA is shown in Fig. 2A. The phylogeny showed that from the three sequences characterized in this study, one clustered with HDV-3 and two with African HDV-8. The HDV-3 sequence obtained herein closely grouped with other HDV-3 sequences isolated in the Amazonas state (Western Amazon Basin).

The two HDV-8 sequences from Maranhão state in Brazil showed 78.8–87.7% (mean 83.5%) similarity with the other ten HDV-8 African sequences previously described and 89.4% with




Fig. 1. Geographic location of Maranhão state in Brazil.

each other. The 19 amino acids of the L-HDAg carboxyl terminus contained some amino acids that were exclusive for the Brazilian HDV-8 sequences (data not shown).

HBV subgenotypes from these cases were also characterized by phylogenetic analyses (Fig. 2B). The two patients infected with HDV-8 had co-infection with HBV subgenotype D4 (ayw2). The sequence that clustered closer with HBV subgenotype D4 sequences (FJ0100650) was from Boca do Acre, Amazonas state. Other sequences in a closer tree branch also clustering with a high posterior probability value were from Rondônia state, which is close to Boca do Acre.

The single patient infected with HDV-3 was co-infected with HBV subgenotype A1 (adw2). This HBV sequence clustered with other Brazilian sequences from different states and it was closer to two sequences from the Amazon region (Rondônia and Amazonas states) and two other sequences obtained in Rio de Janeiro, Southeast Brazil.

#### 4. Discussion

In this study, we identified the presence of HDV infection in Maranhão state, a region not included in the HDV endemic area of

Brazil. A prevalence of 3.8% (5/133) of antibodies against HDV (anti-HD) was found among HBsAg chronic carriers from this region. This prevalence was lower than that found in endemic areas (Amazon region) where the prevalence of anti-HD in HBsAg positive patients reached 34.4% (Fonseca et al., 1988; Viana et al., 2005), but was higher than that already described in regions other than the endemic area (Ferraz et al., 1985; Oliveira et al., 1999; Strauss et al., 1987).

It is noteworthy that most of the anti-HD positive patients found in Maranhão state were from Urbano Santos, a city located in a rural area in the east of the state. These patients were not from the same family and no close relation between them was identified.

The finding of HDV infection in all the three patients from Urbano Santos studied herein emphasizes the need for more careful studies to analyze the epidemiology of hepatitis viruses infection in this area, as our results suggests that this area possibly represents an endemic area for HDV within Maranhão state in Brazil. Additionally, in a detailed medical interview, these patients reported that the occurrence of liver-related complications (especially ascitis and jaundice) has been associated with previous deaths in their families, suggesting that HBV/HDV infection may have been frequent in this area for a long time.


**Table 1**  
Demographic, clinical, and virological features of anti-HD positive patients from Maranhão state, Northeast of Brazil.

Patient	Gender	Age	Race <sup>a</sup>	Origin (city)	HBeAg/anti-HBe status	HBV DNA levels (log)	HDV RNA	HBV subgenotype and subtype	HDV genotype	Clinical status
8127	M	41	Brown	Urbano Santos	–/+	3.43	+	D4 (ayw2)	HDV-8	Cirrhosis
8103	F	61	Brown	Urbano Santos	–/+	4.74	–	ND	ND	Cirrhosis
8135	M	53	Black	São Luís	–/+	2.63	–	ND	ND	Inactive carrier
8022	M	78	Brown	Urbano Santos	+/+	4.91	+	D4 (ayw2)	HDV-8	Cirrhosis
8141	F	50	Brown	São Luís <sup>b</sup>	–/+	2.49	+	A1 (adw2)	HDV-3	Inactive carrier

HBV DNA levels are expressed as log<sub>10</sub> IU/ml; ND: not done.

<sup>a</sup> Based on the skin color.

<sup>b</sup> This patient lives in São Luís, but was born in Manaus, a city located in the Amazonas State, in the Western Amazon Basin, Brazil.



**Fig. 2.** Maximum clade credibility (MCC) trees estimated by Bayesian analysis of (A) 108 partial sequences (105 from GenBank and 3 from this study) of delta antigen gene; (B) 284 sequences (281 from GenBank and 3 from this study) partially comprising HBsAg and DNA polymerase coding regions (S/POL). Sequences characterized in this study are highlighted with a triangle and sequences obtained from GenBank are identified by their accession number followed by an indication of the geographic origin. In (A) the collapsed clades correspond to sequences of HDV genotypes 1, 2, 4, 5, 6, and 7; and in (B) they correspond to sequences of other HBV genotypes (B, C, E, F, G, H, I) and the subgenotypes of genotype A (A2–A7) and D (D1, D2, D3, D5, and D7). The values of posterior probability are shown above their respective key nodes.

A recent population-based study evaluating only the populations from the capital cities of Brazil report that only 25% of the population is vaccinated against HBV in the Northeast region (Pereira et al., 2009). Further studies should be done to better evaluate the percentage of susceptible individuals in the Northeast states and to implement HBV vaccination campaigns in these areas, especially in regions where HDV infection is also reported.

In this study, HDV genotype 3 was found in a patient from São Luís, the capital city of Maranhão State. HDV-3 has been observed exclusively in the northern region of South America (Casey et al., 1993; Gomes-Gouvea et al., 2008, 2009; Quintero et al., 2001). This

patient lives in Maranhão, but she was born in an HDV endemic region (Amazonas state, Western Amazon region). Thus, she possibly contracted HDV infection there, as the HDV-3 sequence obtained from her sample grouped closer to HDV-3 sequences previously isolated in Amazonas state (Gomes-Gouvea et al., 2009). The genotype A (subgenotype A1) of HBV found in this patient is also frequent in the Brazilian Amazon Basin and has also been found in association with HDV-3 (Gomes-Gouvea et al., 2009). The HBV sequence also clustered with other sequences obtained from the Amazonas state, reinforcing the hypothesis that this patient was most likely contaminated in that region.



Interestingly, we found an HDV genotype not found before in South America and only reported in individuals from West Africa. Two patients who were born and have always lived in Urbano Santos County in Maranhão state were infected with HDV-8.

HDV genotype 8 was initially isolated in France from three patients from three different West African countries: Congo, Senegal, and Côte d'Ivoire (Le Gal et al., 2006). Recently, HDV-8 was identified in rural and urban populations from Gabon (Central Africa), reinforcing the African origin of this genotype (Makuwa et al., 2008, 2009).

There are few data on HDV genotype distribution in the African continent. The studies done so far show that HDV-1 is currently the most prevalent genotype, followed by HDV-8, HDV-5, HDV-7, and HDV-6 (Le Gal et al., 2006; Makuwa et al., 2008, 2009; Radjef et al., 2004; Yacouba et al., 2011; Zhang et al., 1996). The new characterized African genotypes (HDV-5 to HDV-8) may represent the ancient genotypes of HDV that were the most prevalent genotypes in the past. Currently, these genotypes are found circulating in limited areas in some regions of Africa for reasons not yet known.

It is estimated that from XVI to XVIII century, millions of Africans were taken to Brazil as slaves (Voyages: The Trans-Atlantic Slave Trade Database. <http://www.slavevoyages.org>). This group, together with Amerindians and Europeans, represent the three major ancestral roots that originated the Brazilian people (Riberiro, 2006). In Brazil, and particularly in the Maranhão state, numerous Quilombos (which are communities formed mostly by descendants of African slaves) exist. We have not found any evidence of a recent relation among our patients and the African continent, thus the results suggest that HDV-8 might have come to Maranhão state a long time ago together with Africans taken to Brazil as slaves.

Interesting, the two cases infected with HDV-8 were also co-infected with HBV subgenotype D4. This subgenotype was found previously in two other regions in northern Brazil (Gomes-Gouveia et al., 2009; Santos et al., 2010) and in Haiti, another country where Africans were massively taken as slaves (Andernach et al., 2009). The African slaves who went to northern Brazil and the Caribbean Islands were mostly from the North Atlantic, and far away from the southern African regions that supplied the rest of Brazil (Domingues da Silva, 2008). Thus, according to historical records, the presence of HBV/D4 in two different regions in Latin America might reflect the origin of slaves and routes of slave traffic ships.

Currently, this subgenotype is not frequent in Africa and has been only described in Ruanda (Hubschen et al., 2009). Nevertheless, it is possible that this genotype was largely distributed in different regions of the African continent in the past, being later replaced by a different genotype.

Herein we showed that HDV infection is present in the North-west region of Brazil. This is worrisome and highlights the need to implement clinical and epidemiological studies to clarify the presence of HDV infection in areas other than the ones already known as endemic.

The presence of HDV-8 infected individuals in Brazil who have not been in Africa may reflect the close relation with HDV genotypes' geographic distribution and human migrations.

## Acknowledgments

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP 2010/50081-9 and 2010/51208-2, Fundação de Amparo à Pesquisa do Estado do Maranhão – FAPEMA (APP-2009/00950) and Secretaria de Saúde do Estado do Maranhão – Departamento de Vigilância Epidemiológica.

## References

- Andernach, I.E., Nolte, C., Pape, J.W., Muller, C.P., 2009. Slave trade and hepatitis B virus genotypes and subgenotypes in Haiti and Africa. *Emerg. Infect. Dis.* 15 (8), 1222–1228.
- Bensabath, G., Dias, L.B., 1983. Labrea hepatitis (Labrea black fever) and other fulminant forms of hepatitis in Sena Madureira, Acre and Boca do Acre, Amazonas, Brazil. *Rev. Inst. Med. Trop. Sao Paulo* 25 (4), 182–194.
- Bensabath, G., Hadler, S.C., Soares, M.C., Fields, H., Dias, L.B., Popper, H., Maynard, J.E., 1987. Hepatitis delta virus infection and Labrea hepatitis. Prevalence and role in fulminant hepatitis in the Amazon Basin. *JAMA* 258 (4), 479–483.
- Casey, J.L., 1996. Hepatitis delta virus. Genetics and pathogenesis. *Clin. Lab. Med.* 16 (2), 451–464.
- Casey, J.L., Brown, T.L., Colan, E.J., Wignall, F.S., Gerin, J.L., 1993. A genotype of hepatitis D virus that occurs in northern South America. *Proc. Natl. Acad. Sci. U.S.A.* 90 (19), 9016–9020.
- Domingues da Silva, D.B., 2008. The Atlantic slave trade to Maranhão, 1680–1846: volume, routes and organisation. *Slavery Abolition* 29 (4), 477–501.
- Drummond, A.J., Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* 7, 214.
- Ferraz, M.L., Vilela, M.P., Silva, A.E., Paiva, E.R., 1985. Serological markers for hepatitis B virus and delta antibody in Brazilian male homosexuals. *Rev. Paul. Med.* 103 (5), 228–230.
- Fonseca, J.C., Simonetti, S.R., Schatzmayr, H.G., Castejon, M.J., Cesario, A.L., Simonetti, J.P., 1988. Prevalence of infection with hepatitis delta virus (HDV) among carriers of hepatitis B surface antigen in Amazonas State, Brazil. *Trans. Roy. Soc. Trop. Med. Hyg.* 82 (3), 469–471.
- Gomes-Gouveia, M.S., Pereira Soares Mdo, C., Guedes de Carvalho Mello, I.M., Brito, E.M., Pereira Moia Lde, J., Bensabath, G., Nunes, H.M., Carrilho, F.J., Pinho, J.R., 2008. Hepatitis D and B virus genotypes in chronically infected patients from the Eastern Amazon Basin. *Acta Trop.* 106 (3), 149–155.
- Gomes-Gouveia, M.S., Soares, M.C., Bensabath, G., de Carvalho-Mello, I.M., Brito, E.M., Souza, O.S., Queiroz, A.T., Carrilho, F.J., Pinho, J.R., 2009. Hepatitis B virus and hepatitis delta virus genotypes in outbreaks of fulminant hepatitis (Labrea black fever) in the western Brazilian Amazon region. *J. Gen. Virol.* 90 (Pt 11), 2638–2643.
- Hubschen, J.M., Mugabo, J., Peltier, C.A., Karasi, J.C., Sausy, A., Kirpach, P., Arendt, V., Muller, C.P., 2009. Exceptional genetic variability of hepatitis B virus indicates that Rwanda is east of an emerging African genotype E/A1 divide. *J. Med. Virol.* 81 (3), 435–440.
- Ivaniushina, V., Radjef, N., Alexeeva, M., Gault, E., Semenov, S., Salhi, M., Kiselev, O., Deny, P., 2001. Hepatitis delta virus genotypes I and II cocirculate in an endemic area of Yakutia, Russia. *J. Gen. Virol.* 2 (Pt 11), 2709–2718.
- Kwok, S., Higuchi, R., 1989. Avoiding false positives with PCR. *Nature* 339 (6221), 237–238.
- Le Gal, F., Gault, E., Ripault, M.P., Serpaggi, J., Trinchet, J.C., Gordien, E., Deny, P., 2006. Eighth major clade for hepatitis delta virus. *Emerg. Infect. Dis.* 12 (9), 1447–1450.
- Lee, C.M., Changchien, C.S., Chung, J.C., Liaw, Y.F., 1996. Characterization of a new genotype II hepatitis delta virus from Taiwan. *J. Med. Virol.* 49 (2), 145–154.
- Makuwa, M., Caron, M., Souquiere, S., Malonga-Moulet, G., Mahe, A., Kazanji, M., 2008. Prevalence and genetic diversity of hepatitis B and delta viruses in pregnant women in Gabon: molecular evidence that hepatitis delta virus clade 8 originates from and is endemic in central Africa. *J. Clin. Microbiol.* 46 (2), 754–756.
- Makuwa, M., Mints-Ndong, A., Souquiere, S., Nkoghe, D., Leroy, E.M., Kazanji, M., 2009. Prevalence and molecular diversity of hepatitis B virus and hepatitis delta virus in urban and rural populations in northern Gabon in central Africa. *J. Clin. Microbiol.* 47 (7), 2265–2268.
- Oliveira, M.L., Bastos, F.L., Telles, P.R., Yoshida, C.F., Schatzmayr, H.G., Paetzold, U., Pauli, G., Schreier, E., 1999. Prevalence and risk factors for HBV, HCV and HDV infections among injecting drug users from Rio de Janeiro, Brazil. *Braz. J. Med. Biol. Res.* 32 (9), 1107–1114.
- Pereira, L.M., Martelli, C.M., Merchan-Hamann, E., Montarroyos, U.R., Braga, M.C., de Lima, M.L., Cardoso, M.R., Turchi, M.D., Costa, M.A., de Alencar, L.C., Moreira, R.C., Figueiredo, G.M., Ximenes, R.A., 2009. Population-based multicentric survey of hepatitis B infection and risk factor differences among three regions in Brazil. *Am. J. Trop. Med. Hyg.* 81 (2), 240–247.
- Posada, D., Crandall, K.A., 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14, 817–818.
- Quintero, A., Uzcatégui, N., Loureiro, C.L., Villegas, L., Illarramendi, X., Guevara, M.E., Ludert, J.E., Bliz, L., Liprandi, F., Pujol, F.H., 2001. Hepatitis delta virus genotypes I and III circulate associated with hepatitis B virus genotype F in Venezuela. *J. Med. Virol.* 64 (3), 356–359.
- Radjef, N., Gordien, E., Ivaniushina, V., Gault, E., Anais, P., Drugan, T., Trinchet, J.C., Roulot, D., Tamby, M., Milinkovitch, M.C., Deny, P., 2004. Molecular phylogenetic analyses indicate a wide and ancient radiation of African hepatitis delta virus, suggesting a deltavirus genus of at least seven major clades. *J. Virol.* 78 (5), 2537–2544.
- Ribeiro, D., 2006. O povo Brasileiro: A formação e o sentido do Brasil, 1st ed. Companhia das Letras, São Paulo.
- Rizzetto, M., 2009. Hepatitis D: thirty years after. *J. Hepatol.* 50 (5), 1043–1050.
- Sakugawa, H., Nakasone, H., Nakayoshi, T., Kawakami, Y., Miyazato, S., Kinjo, F., Saito, A., Ma, S.P., Hotta, H., Kinoshita, M., 1999. Hepatitis delta virus genotype IIb predominates in an endemic area, Okinawa, Japan. *J. Med. Virol.* 58 (4), 366–372.

- Santos, A.O., Alvarado-Mora, M.V., Botelho, L., Vieira, D.S., Pinho, J.R., Carrilho, F.J., Honda, E.R., Salcedo, J.M., 2010. Characterization of hepatitis B virus (HBV) genotypes in patients from Rondonia, Brazil. *Viol. J.* 7, 315.
- Shakil, A.O., Hadziyannis, S., Hoofnagle, J.H., Di Bisceglie, A.M., Gerin, J.L., Casey, J.L., 1997. Geographic distribution and genetic variability of hepatitis delta virus genotype I. *Virology* 234 (1), 160–167.
- Sitnik, R., Paes, A., Manguiera, C.P., Pinho, J.R., 2010. A real-time quantitative assay for hepatitis B DNA virus (HBV) developed to detect all HBV genotypes. *Rev. Inst. Med. Trop. Sao Paulo* 52 (3), 119–124.
- Strauss, E., Gayotto, L.C., da Silva, L.C., Alves, V.A., Carrilho, F., Chamone, D.A., da Silva, E.F., Porta, G., Granato, C., Trepo, C., 1987. Unexpected low prevalence of delta antibodies in the east Amazon region and Sao Paulo: evidence for regional differences in the epidemiology of delta hepatitis virus within Brazil. *Trans. Roy. Soc. Trop. Med. Hyg.* 81 (1), 73–74.
- Viana, S., Parana, R., Moreira, R.C., Compri, A.P., Macedo, V., 2005. High prevalence of hepatitis B virus and hepatitis D virus in the western Brazilian Amazon. *Am. J. Trop. Med. Hyg.* 73, 808–814.
- Wedemeyer, H., Manns, M.P., 2010. Epidemiology, pathogenesis and management of hepatitis D: update and challenges ahead. *Nat. Rev. Gastroenterol. Hepatol.* 7 (1), 31–40.
- Yacouba, F., Dominique, N.N., Michele, T.S., Richard, N., 2011. High prevalence and predominance of Hepatitis D genotype 1 infection in Cameroon. *J. Clin. Microbiol.*
- Zhang, Y.Y., Tsega, E., Hansson, B.G., 1996. Phylogenetic analysis of hepatitis D viruses indicating a new genotype I subgroup among African isolates. *J. Clin. Microbiol.* 34 (12), 3023–3030.